

Science & Healthcare


PEER-REVIEWED MEDICAL SCIENTIFIC JOURNAL

**РЕЦЕНЗИРУЕМЫЙ МЕДИЦИНСКИЙ
НАУЧНО-ПРАКТИЧЕСКИЙ ЖУРНАЛ**


Наука и Здравоохранение

Ғылым және Денсаулық Сақтау

**РЕЦЕНЗИЯЛАНАТЫН МЕДИЦИНАЛЫҚ
ҒЫЛЫМИ-ПРАКТИКАЛЫҚ ЖУРНАЛ**



**Volume 26,
Issue 5
2024**



Министерство здравоохранения
Республики Казахстан
Учредитель: НАО
«Медицинский университет Семей»
Основан в 1999 году.

Журнал зарегистрирован в
Министерстве информации и
коммуникаций Республики Казахстан
Комитете информации № 17773-Ж.

Входит в Перечень изданий,
рекомендуемых Комитетом по
обеспечению качества в сфере науки и
высшего образования МНВО
Республики Казахстан для публикации
основных результатов научной
деятельности (Приказ №603 от
12.07.2024г.)

Включен в Ulrich's Periodicals
Directory, Global Health, CAB
Abstracts, InfoBase Index, Directory
of Research Journals Indexing,
Российский индекс научного
цитирования (РИНЦ), E-library.ru,
Cyberleninka.ru, Norwegian register
for scientific journals (NSD),
Всесоюзный институт научной и
технической информации Российс-
кой академии наук (ВИНИТИ РАН),
Ассоциацию CONEM, РАЦС,
DataBase Indexing, ICI World of
Journals

Подписной индекс 74611
в каталоге «Казпочта»
Цена свободная

Сайт <http://newjournal.ssmu.kz>
e-mail: selnura@mail.ru

Адрес редакции:
071400, г. Семей
ул. Абая Кунанбаева, 103
контактный телефон:
(7222) 56-42-09 (вн. № 1054)
факс: (7222) 56-97-55

Выпускающий редактор:
Э.Ф. Сапаргалиева

Переводчики:
С.А. Жаукенова, Н.А. Шумский



При цитировании материалов
ссылка на журнал обязательна.
Отпечатано в типографии
Медицинского университета Семей
Подписано в печать: 30.10.2024г.
Формат 60x90/8.
Печать цифровая. Усл.п.л 30,8
Тираж 500 экз., зак.184

Online ISSN: 3007 – 0244
Print ISSN: 2410 – 4280

НАУКА И ЗДРАВООХРАНЕНИЕ

РЕЦЕНЗИРУЕМЫЙ МЕДИЦИНСКИЙ НАУЧНО-ПРАКТИЧЕСКИЙ ЖУРНАЛ

Том 26 №5, 2024

Журнал «Наука и Здравоохранение» - рецензируемый медицинский научно-практический журнал, который публикует результаты оригинальных исследований, обзоры литературы, клинические случаи, краткие сообщения и отчеты о конференциях по широкому кругу вопросов, связанных с клинической медициной и общественным здоровьем. Основной читательской аудиторией журнала является медицинское научное сообщество, практикующие врачи, докторанты и магистранты в области медицины и общественного здоровья.

Главный редактор: **А.А. Дюсупов**
доктор медицинских наук, профессор

Зам. главного редактора: **Д.М. Шабдарбаева**
доктор медицинских наук, профессор

Редакционный совет:

Абдрахманов А.С. (Нур-Султан, Казахстан)
Акильжанова А.Р. (Нур-Султан, Казахстан)
Акшулаков С.К. (Нур-Султан, Казахстан)
Баймаханов Б.Б. (Алматы, Казахстан)
Брузати Лука Джиованни Карло (Удин, Италия)
Виджай Кумар Чатту (Торонто, Канада)
Гржибовский А.М. (Архангельск, Российская Федерация)
Гюрель Фазыл Сердар (Анкара, Турция)
Даутов Т.Б. (Нур-Султан, Казахстан)
Джерзи Крупински Белецки (Барселона, Испания)
Жумадилов Ж.Ш. (Нур-Султан, Казахстан)
Кавальчи Чемиль (Анкара, Турция)
Карпенко А.А. (Новосибирск, Российская Федерация)
Ковальчук В.В. (Санкт-Петербург, Российская Федерация)
Лесовой В.Н. (Харьков, Украина)
Лукьянов С.А. (Москва, Российская Федерация)
Мутиг К. (Шарите, Германия)
Носо Й. (Шимане, Япония)
Раманкулов Е.М. (Нур-Султан, Казахстан)
Степаненко В.Ф. (Обнинск, Российская Федерация)
Тапбергенов С.О. (Семей, Казахстан)
Тринчеро Элизабетта Флора Ольга (Милан, Италия)
Хоссейни Хенгаме (Скрантон, Соединенные Штаты Америки)
Хоши М. (Хиросима, Япония)
Шейнин А. (Тель-Авив, Израиль)

Содержание

ОРИГИНАЛЬНЫЕ ИССЛЕДОВАНИЯ

- Zhautikova S.B., Abikenova F.S., Zhiembayeva K.M., Bakaramova G.A., Kim V., Volodina A., Dosanova A.** 7-15
Analysis of correlation and regression relationships between thyroid dysfunction, adrenal dysfunction and cardiopulmonary disorders in patients with diabetes mellitus
- Казарян С.У., Базарбекова Р.Б., Джунусбекова Г.А., Досанова А.К., Кайнарбекова Н.Б.** 16-24
Состояние коронарного русла у пациентов с острым коронарным синдромом на фоне впервые выявленных нарушений углеводного обмена
- Rakhmankulova A.M., Pak L.A., Romyantsev P.O., Pivina L.M., Esbolatova N.S., Bolsynbekova S.O., Seitkhanova D.M., Bakhtiyar A.S., Baurzhan A., Zharlyganova D.S.** 25-32
The clinical features of radioiodine-resistant differentiated thyroid cancer in patients of the republic of Kazakhstan: preliminary results
- Оразбаева Д.Р., Кенсарина А.Н., Орындыбаева А.К., Ахметов Е.А., Манатова А.М., Куаныш Ж.М., Жарлыганова Д.С.** 33-40
Анализ результатов ультразвуковой диагностики с доплерографией при доброкачественных и злокачественных опухолях щитовидной железы
- Еспенбетова М.Ж., Бидакхметова А.М., Баймуханова А.А., Крыкпаева А.С., Уашева Л.Б.** 41-46
Частота встречаемости узловых форм щитовидной железы по Семейскому региону
- Turmanbayeva A.A., Sadykova K.Zh., Nuskabayeva G.O., Nurdinov N.S., Sarsenbayeva G.Zh., Kemelbekov K.S., Seidakhmetova A.A.** 47-54
The relationship between the main lifestyle factors and metabolic syndrome
- Bekenova N.B., Aitkaliev A.D., Kassiyeva B.S., Vochshenkova T.A., Suleimenova A.B., Benberin V.V.** 55-61
Relationship of the rs564398 polymorphism of the ANRIL gene (CDKN2B) with complications of type 2 diabetes, particularly with the cardiovascular form of diabetic neuropathy
- Sarshayev M.A., Berdikhojayev M.S., Karibayev K.R., Turdaliyeva B.S., Tanbayeva G.Z., Makhanbetkhan S.S., Biniyazova A.A., Lupezhova A.D., Zhumabekov A.K.** 62-71
Risk factors for mortality after stenting of symptomatic atherosclerotic stenoses in intra- and extracranial arteries: a cross-sectional study
- Kussainova Zh., Bulegenov T., Zhapar Zh., Abiralina Sh., Abiltayev A., Tashtemirova O., Abilmazhinova G.** 72-78
Organization of preventive measures for infective endocarditis in Kazakhstan: a cross-sectional study among cardiologists

Table Of Contents

ORIGINAL ARTICLES

- Жаутикова С.Б., Абикенова Ф.С., Жиенбаева К.М., Бакарамова Г.А., Ким В., Володина А., Досанова А.**
Анализ корреляционных и регрессионных связей между нарушением функции щитовидной железы, надпочечников и кардиопульмональными нарушениями у больных сахарным диабетом
- Kazaryan S.U., Bazarbekova R.B., Dzhunusbekova G.A., Dosanova A.K., Kainarbekova N.B.**
Condition of coronary arteries in patients with acute coronary syndrome caused by newly identified disorders of carbohydrate metabolism
- Рахманкулова А.М., Пак Л.А., Румянцев П.О., Пивина Л.М., Есболатова Н.С., Болсынбекова С.О., Сейтханова Д.М., Бактияр А.С., Бауржан А.Б., Жарлыганова Д.С.**
Клинические особенности радиойодрезистентного дифференцированного рака щитовидной железы у пациентов Республики Казахстан: предварительные результаты
- Orazbaeva D.R., Kensarina A.N., Oryndybaeva A.K., Akhmetov E.A., Manatova A.M., Kuanysh Zh.M., Zharlyganova D.S.**
Analysis of the results of ultrasound diagnostics with Dopplerography for benign and malignant tumors of the thyroid gland
- Espenbetova M.Zh., Bidakhmetova A.M., Baimukhanova A.A., Krykpaeva A.S., Uasheva L.B.**
Prevalence of Nodular Thyroid Forms in the Semey Region
- Турманбаева А.А., Садыкова К.Ж., Нускабаева Г.О., Нурдинов Н.С., Сарсенбаева Г.Ж., Кемельбеков К.С., Сейдахметова А.А.**
Взаимосвязь между основными факторами образа жизни и метаболическим синдромом
- Бекенова Н.Б., Айткалиев А.Д., Касиева Б.С., Вощенко Т.А., Сулейменова А.Б., Бенберин В.В.**
Связь полиморфизма rs564398 гена ANRIL (CDKN2B) с осложнениями сахарного диабета 2 типа, в частности с сердечно-сосудистой формой диабетической нейропатии
- Саршаев М.А., Бердиходжаев М.С., Карибаев К.Р., Турдалиева Б.С., Танбаева Г.З., Маханбетхан Ш.С., Биниязова А.А., Лупежова А.Д., Жумабеков А.К.**
Факторы риска смертности после стентирования симптоматических атеросклеротических стенозов в интра- и экстракраниальных артериях: поперечное исследование
- Құсәинова Ж., Булегенов Т., Жапар Ж., Абралина Ш., Абильтәев А., Таштемирова О., Абильтәминова Г.**
Организация профилактических мер при инфекционном эндокардите в Казахстане: поперечное исследование среди кардиологов

Оразбаева Д.Р., Иса Г., Сагындыков И.К., Ахметов Е.А., Манатова А.М., Жарлыганова Д.С.
Роль ультразвукового исследования в выявлении сосудистых осложнений у реципиентов трансплантата печени

Skenderova A.A., Nuskabayeva G.N., Tatykayeva U.B., Sarsenbayeva G.Zh., Kemelbekov K.S., Seidakhmetova A.A.
Therapeutic effects of melatonin in an ethanol-induced gastric ulcer model

Bedel H.-A., Bedel C., Selvi F., Zortuk Ö., Karanci Yu.
Effect of Seasonal Changes on scorpion poison
Абуова А.С., Мейрамова Ә.М., Абдахина Б.Б., Бокаева А.А., Цой Т.Т., Бикташев Д.Б., Айнабекова Б.А.

Особенности течения анкилозирующего спондилоартрита в казахстанской популяции в зависимости от гендерной принадлежности
Apbassova M., Apbassova S.A., Shabdarbayeva D.M., Chaizhunusova N.Zh., Bulegenov T.A., Mutig K., Mussina A.A., Baurzhan A.B., Assubayev O.B., Dyusembayev A.K., Lepesbayev M.N., Amantayeva G.K., Tatykayeva U.B., Svitich T.N., Shalgumbayeva G.M., Dyussupov A.A.
Suicide rate forecasting in the cities of Ust-Kamenogorsk, Semey, and Pavlodar region using a mathematical model

Byltash S., Brimzhanova M., Faizullina K., Akhtayeva N., Alekenova N., Kosherbayeva L., Nurbaulina E.
Current practice in caring for people with dementia
Akylzhanova Zh.E., Tanysheva G.A., Gasanova E.Z., Bilyalova G.T., Nurmagambetova D.M., Akhmetova A.B., Suleimenova A.P., Ulan S.M., Nurmakhan N.S.
Using the Robson classification to estimate the caesarean section rate and look for ways to reduce it

COVID-19 - АКТУАЛЬНАЯ ТЕМА

Дюсупов А.А., Жуманбаева Ж.М., Кайрханова Ы.О., Досбаева А.Д., Крыкпаева А.С., Серикбаев А.С., Амренова К.Ш., Сеиткабылов А.А., Кудайбергенов Р.Н.
Пост-COVID-19-синдром: симптомы и факторы риска

ОБЗОР ЛИТЕРАТУРЫ

Shokenov R.D., Shakhmarova T.K., Mirmanova Zh.Zh., Chamoieva A.Ye., Zhalbinova M.R., Rakhimova S.E., Bekbossynova M.S., Akilzhanova A.R.
Characterizing the Role of ABCG5/G8 in Sitosterolemia: Diagnostic Challenges in Differentiating from Familial Hypercholesterolemia
Samatova N.E., Kamalbekova G.M., Jaxalykova K.K., Mussina A.A., Karin B.T.
The main tools of the confidential audit of perinatal mortality: a review

79-84 Orazbaeva D.R., Isa G., Sagyndykov I.K., Akhmetov E.A., Manatova A.M., Zharlyganova D.S.
The role of ultrasound in identifying vascular complications in liver transplant recipients

85-92 Скендерова А.А., Нускабаева Г.О., Татыкаева У.Б., Сарсенбаева Г.Ж., Кемельбеков К.С., Сейдахметова А.А.
Терапевтические эффекты мелатонина на модели язвы желудка, вызванной этанолом

93-98 Бедел Х.-А., Бедел Дж., Сельви Ф., Зортук О., Каранджи Ю.
Влияние сезонных изменений на яд скорпиона

99-107 Abuova A.S., Meiramova A.M., Abdakhina B.B., Bokayeva A.A., Tsoy T.T., Biktashev D.B., Ainabekova B.A.
Features of the ankylosing spondylitis course depending on gender in the Kazakh population

108-114 Апбасова М., Апбасова С.А., Шабдарбаева Д.М., Чайжунусова Н.Ж., Булегенов Т.А., Мутиг К., Мусина А.А., Бауржан А.Б., Асубаев О.Б., Дюсембаев А.К., Лепесбаев М.Н., Амантаева Г.К., Татыкаева У.Б., Свитич Т.Н., Шалгумбаева Г.М., Дюсупов А.А.
Прогнозирование уровня суицида в городах Усть-Каменогорск, Семей и Павлодарской области с помощью математической модели

115-120 Былташ С., Бримжанова М., Файзуллина К., Ахтаева Н., Алеkenова Н., Кошербаева Л., Нурбаулина Э.
Существующая практика ухода за людьми с деменцией

121-128 Акылжанова Ж.Е., Танышева Г.А., Гасанова Э.З., Билялова Г.Т., Нурмагамбетова Д.М., Ахметова А.Б., Сулейменова Ә.С., Ұлан С.М., Нурмахан Н.С.
Применение классификации Робсона для оценки показателя частоты кесарева сечения и поиска возможностей его снижения

COVID-19 - TOPICAL SUBJECT

129-140 Dyussupov A.A., Zhumanbayeva Zh.M., Kairkhanova Y.O., Dosbayeva A.M., Krykpaeva A.S., Serikbayev A.S., Amrenova K.Sh., Seitkabylov A.A., Kudaibergenov R.N.
Post-COVID-19 syndrome: symptoms and risk factors

REVIEWS

141-149 Шокенов Р.Д., Шахмарова Т.К., Мирманова Ж.Ж., Чамойева А.Е., Жалбинова М.Р., Рахимова С.Е., Бекбосынова М.С., Акильжанова А.Р.
Характеристика роли ABCG5/G8 при ситостеролемии: диагностические трудности в дифференциации от семейной гиперхолестеринемии

150-159 Саматова Н.Е., Камалбекова Г.М., Джаксалыкова К.К., Мусина А.А., Карин Б.Т.
Основные инструменты конфиденциального аудита перинатальной смертности: обзор литературы

Ракишева Ж.В.

Актуальное состояние проблемы нарушения развития нервной системы в раннем возрасте и последующее вмешательство. Обзор литературы

Kussainova A.A., Adilzhanov A.B., Adilgozhin M.S., Shalgumbayeva G.M., Derbissalina G.A., Bekbergenova Zh.B., Umbetzhanova A.T., Kozhakhmetova D.K., Zhokebaeva M.S., Shamshudinov T.M., Kassym L.T.

Tuberculosis: prevalence, risk factors, and first-line drug resistance

Ibraeva A.K., Latypova N.A., Kushugulova A.R., Meiramova A.M., Adilbekova B.B., Isilbaeva A.A., Kozhakhmetov S.S.

Probiotics as additional treatment of Rheumatoid arthritis. A review

Канапиянов Б., Нуртазина А., Даутов Д., Тойшиманов М., Войцеховский И., Нургалиева Б.

Распространенность предиабета в азиатской популяции: Обзор литературы

Espenbetova M.Zh., Bidakhmetova A.M., Krykpaeva A.S., Uasheva L.B., Syzdykbaev M.K., Baimukhanova A.A., Baurzhan A.B.

Prognostic Significance of Genetic Tumor Markers in Thyroid Neoplasms: Literature Review

КЛИНИЧЕСКИЙ СЛУЧАЙ

Абдикаримов А.М., Омаров Е.Д., Сабербеков С.О., Мукантаев Е.Т., Ерпашов А.П., Төлегенұлы А.

Первый опыт и успешное хирургическое лечение рака левой почки с протяженным опухолевым венозным тромбозом нижней полой вены и правого предсердия [IV уровня] в Республике Казахстан

Tynyshtykbayeva N.Ye., Aitkulova A.M., Kadenova T.B., Rakhimova S.Ye., Samatkyzy D., Rogounovitch T.I., Kukanova A.M., Begimbetova D.A., Yermekbayeva B.A., Kozhamkulov U.A., Yerezhepov D.A., Akilzhanova A.R., Sarbassov D.D.

Clinical relevance of KRAS mutation testing in metastatic colorectal using different sequencing platforms: a case report

Рустамова Ф.Е., Мухамбетова Р.У., Иминова Ш.И., Баймолда А.Е., Әбдіхалық Н.А., Сқақбаев Ш.Е.

Описание клинического случая генерализованной лимфаденопатии неясной этиологии

Akylzhanova Zh.E., Tanysheva G.A., Тультаев Р.Б., Bilyalova G.T., Nurmagambetova D.M., Шаймуханова З.С., Турсынбаева А.М.

Embolization of the uterine arteries for late postpartum hemorrhage due to metroendometritis: a clinical case

160-176 Rakisheva Zh.V.

The current state of the problem of impaired development of the nervous system at an early age and subsequent intervention. Review

177-187 Кусаинова А.А., Адилжанов А.Б., Адильгожин М.С., Шалгумбаева Г.М., Дербисалина Г.А., Бекбергенова Ж.Б., Умбетжанова А.Т., Кожакметова Д.К., Жокебаева М.С., Шамшудинов Т.М., Касым Л.Т.

Туберкулез: распространенность, факторы риска и устойчивость к препаратам первой линии

188-197 Ибраева А.К., Латыпова Н.А., Кушугулова А.Р., Мейрамова А.М., Адильбекова Б.Б., Исильбаева А.А., Кожакметов С.С.

Пробиотики как дополнительное лечение ревматоидного артрита. Обзор литературы

198-203 Kanapiyanov B., Nurtazina A., Dautov D., Toishimanov M., Voitsekhovskiy I., Nurgaliyeva B.

Prevalence of prediabetes in the Asian population: Literature review

204-212 Еспенбетова М.Ж., Бидакметова А.М., Крыкпаева А.С., Уашева Л.Б., Сыздыкбаев М.К., Баймуханова А.А., Бауржан А.Б.

Прогностическая значимость генетических онкомаркеров новообразований щитовидной железы: обзор литературы

CLINICAL CASE

213-220 Abdikarimov A.M., Omarov Y.D., Saberbekov S.O., Mukantayev Y.T., Yerpashov A.P., Tolegenuly A.

First experience and successful surgical treatment of left kidney cancer with extended tumor venous thrombosis of the inferior vena cavity and right atrium [level IV] in the Republic of Kazakhstan

221-229 Тыныштыкбаева Н.Е., Айткулова А.М., Каденова Т.Б., Рахимова С.Е., Саматкызы Д., Рогунович Т.И., Куканова А.М., Бегимбетова Д.А., Ермекбаева Б.А., Кожамкулов У.А., Ережепов Д.А., Акильжанова А.Р., Сарбасов Д.Д.

Клиническая значимость тестирования мутаций KRAS при метастатическом колоректальном раке с использованием различных платформ секвенирования: клинический случай

230-235 Rustamova F.Y., Mukhambetova R.U., Iminova Sh.I., Baimolda A.Y., Abdykhalyk N.A., Skakbayev Sh.Y.

Description of a clinical case of generalized lymphadenopathy of unknown etiology

236-242 Акылжанова Ж.Е., Танышева Г.А., Тультаев Р.Б., Билялова Г.Т., Нурмагамбетова Д.М., Шаймуханова З.С., Турсынбаева А.М.

Эмболизация маточных артерий при позднем послеродовом кровотечении на фоне метроэндометрита: клинический случай

Received: 08 June 2024 / Accepted: 14 October 2024 / Published online: 31 October 2024

DOI 10.34689/SH.2024.26.5.006

UDC 616.43-056.52



THE RELATIONSHIP BETWEEN THE MAIN LIFESTYLE FACTORS AND METABOLIC SYNDROME

Ainur A. Turmanbayeva¹, <https://orcid.org/0009-0004-4263-8157>**Karlygash Zh. Sadykova**¹, <https://orcid.org/0000-0002-9120-8565>**Gulnaz O. Nuskabayeva**¹, <https://orcid.org/0000-0003-2139-3221>**Nursultan S. Nurdinov**¹, <https://orcid.org/0000-0001-5341-7211>**Gulzat Zh. Sarsenbayeva**², <https://orcid.org/0000-0002-1518-6528>**Kanatghan S. Kemelbekov**², <https://orcid.org/0009-0008-1619-6278>**Aizat A. Seidakhmetova**², <https://orcid.org/0000-0002-5794-3814>¹ Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkestan, Republic of Kazakhstan;² South Kazakhstan Medical Academy, Shymkent, Republic of Kazakhstan.

Abstract

Introduction. Nowadays, metabolic syndrome continues to be a significant challenge in medicine. Experts from the World Health Organization have called metabolic syndrome (MetS) “the pandemic of the 21st century.” MetS is one of the cardiometabolic risk factors with a high prevalence in the adult population and a high cost to public health systems worldwide. The prevalence of MetS is very high: according to WHO, the incidence of MetS among adults is 20-40% and may increase by 50% in the next 20 years. The most threatening complications closely related to metabolic syndrome are cardiovascular diseases and type 2 diabetes mellitus. Due to the high prevalence of metabolic syndrome, its early detection is of great importance for the timely initiation of the prevention of complications.

Aim. To study the association between the main lifestyle and socio-demographic factors and the risk of developing metabolic syndrome.

Material's and method. The present study is part of a large metabolic syndrome research project in Turkestan city (AP19676909) “Investigation of genetic aspects of nutrition, lifestyle and compliance of patients with metabolic syndrome in a 10-year prospective study”. The sample consisted of 552 patients from the attached population of the outpatient clinic of the H.A. Yasavi International Kazakh-Turkish University. Individuals with acute conditions requiring emergency care, chronic severe decompensated conditions were not included in the study.

Results: The final patient sample was 549 patients, as study participants without data on any of the sociodemographic factors and any of the lipid and carbohydrate metabolism parameters were excluded from the analysis. The association between the presence of MetS and lifestyle indicators, including time spent sitting during the working day and physical activity, was assessed using multivariate logistic regression analysis. After adjusting for the potential confounding factors, such as gender, smoking, and alcohol consumption, the association between MetS and the sitting time during the working day was maintained statistical significance. In contrast, the adjusted logistic regression analysis revealed a statistically significant inverse relationship between the metabolic syndrome and level of physical activity.

Conclusion: The results of our study suggest a significant role for modifiable risk factors, such as a sedentary lifestyle and the level of physical activity, in the development of metabolic syndrome among our study participants. Personalized management tactics for patients with metabolic syndrome, by correcting modifiable lifestyle factors, can help prevent complications.

Keywords: metabolic syndrome, physical activity, lifestyle, sedentary behavior.

Резюме

ВЗАИМОСВЯЗЬ МЕЖДУ ОСНОВНЫМИ ФАКТОРАМИ ОБРАЗА ЖИЗНИ И МЕТАБОЛИЧЕСКИМ СИНДРОМОМ

Айнур А. Турманбаева¹, <https://orcid.org/0009-0004-4263-8157>**Карлыгаш Ж. Садыкова**¹, <https://orcid.org/0000-0002-9120-8565>**Гульназ О. Нускабаева**¹, <https://orcid.org/0000-0003-2139-3221>**Нурсултан С. Нурдинов**¹, <https://orcid.org/0000-0001-5341-7211>**Гульзат Ж. Сарсенбаева**², <https://orcid.org/0000-0002-1518-6528>**Канатжан С. Кемельбеков**², <https://orcid.org/0009-0008-1619-6278>**Айзат А. Сейдахметова**², <https://orcid.org/0000-0002-5794-3814>¹ Международный казахско-турецкий университет имени Ходжи Ахмеда Ясави, г. Туркестан, Республика Казахстан;² АО «Южно-Казахстанская медицинская академия», г. Шымкент, Республика Казахстан.

Актуальность. На сегодняшний день метаболический синдром остается актуальной проблемой медицины. Эксперты Всемирной организации здравоохранения назвали метаболический синдром (МС) «пандемией XXI века». Метаболический синдром (МС) является одним из кардиометаболических факторов риска, имеющим высокую распространенность среди взрослого населения и высокую стоимость для систем общественного здравоохранения во всем мире. По данным ВОЗ частота МС среди взрослого населения составляет 20-40% и может увеличиться на 50% в ближайшие 20 лет. Наиболее грозными осложнениями, тесно связанными с метаболическим синдромом, являются сердечно-сосудистые заболевания и сахарный диабет 2 типа. В связи с высокой распространенностью метаболического синдрома его раннее выявление имеет большое значение для своевременной профилактики осложнений.

Цель: Изучить связь между основными факторами образа жизни и социально-демографическими факторами с риском развития метаболического синдрома.

Материалы и методы: Настоящее исследование является частью научного проекта по изучению метаболического синдрома в городе Туркестан «Изучение генетических аспектов питания, образа жизни и комплаентности пациентов с метаболическим синдромом в 10-летнем проспективном исследовании» (AP19676909). Выборку составили 552 пациента из прикрепленного населения поликлиники Международного казахско-турецкого университета имени Х.А. Ясауи. В исследование не включались пациенты с острыми состояниями, требующими неотложной помощи, хроническими тяжелыми декомпенсированными состояниями.

Результаты: Окончательная выборка пациентов составила 549 человек, так как участники, у которых отсутствовали данные по одному из социально-демографических факторов, либо по одному из параметров липидного и углеводного обмена были исключены из статистического анализа. Связь между наличием метаболического синдрома и показателями образа жизни, включая время, проведенное сидя в течение рабочего дня, физическую активность, была оценена с помощью многомерного логистического регрессионного анализа. Выявлена положительная ассоциация между метаболическим синдромом и временем, проведенным сидя, которая сохраняла статистическую значимость после коррекции на потенциальные конфаундинг факторы, такие как пол, курение и употребление алкоголя. Напротив, в результате скорректированного регрессионного анализа была обнаружена статистически значимая обратная связь метаболического синдрома с уровнем физической активности.

Выводы: Результаты исследования указывают на значительную роль модифицируемых факторов риска (малоподвижный образ жизни, уровень физической активности) в развитии метаболического синдрома в нашей выборке. Персонализированная тактика ведения пациентов с метаболическим синдромом путем коррекции модифицируемых факторов образа жизни может помочь предотвратить осложнения.

Ключевые слова: метаболический синдром, физическая активность, образ жизни, малоподвижный образ жизни.

Түйіндеме

ӨМІР САЛТЫНЫҢ НЕГІЗГІ ФАКТОРЛАРЫ МЕН ЗАТ АЛМАСУ СИНДРОМЫНЫҢ ӨЗАРА БАЙЛАНЫСЫ

Айнур А. Турманбаева¹, <https://orcid.org/0009-0004-4263-8157>

Карлыгаш Ж. Садыкова¹, <https://orcid.org/0000-0002-9120-8565>

Гульназ О. Нускабаева¹, <https://orcid.org/0000-0003-2139-3221>

Нурсултан С. Нурдинов¹, <https://orcid.org/0000-0001-5341-7211>

Гульзат Ж. Сарсенбаева², <https://orcid.org/0000-0002-1518-6528>

Канатжан С. Кемельбеков², <https://orcid.org/0009-0008-1619-6278>

Айзат А. Сейдахметова², <https://orcid.org/0000-0002-5794-3814>

¹ Қ.А. Ясауи атындағы Халықаралық қазақ-түрік университеті, Түркістан қ., Қазақстан Республикасы;

² «Оңтүстік Қазақстан медицина академиясы» АҚ, Шымкент қ., Қазақстан Республикасы.

Өзектілігі. Қазіргі таңда медицинада метаболизмдік синдром өзекті мәселе болып табылады. Дүниежүзілік денсаулық сақтау ұйымының сарапшылары метаболизмдік синдромды «XXI ғасырдың пандемиясы» деп сипаттады. Метаболизмдік синдром - ересек популяциялар арасында таралуы жоғары және бүкіл әлем бойынша қоғамдық денсаулық сақтау жүйелері үшін жоғары шығындары бар кардиометаболикалық қауіп факторларының бірі. ДДҰ деректері бойынша ересек популяцияда МС жиілігі 20-40% құрайды, ал келесі 20 жылда ол 50% артуы мүмкін. Аталған синдроммен тығыз байланысты асқынулары жүрек-қан тамырлары аурулары және 2 типті қант диабеті болып табылады. Метаболизмдік синдромның жоғары жиілікте таралуына байланысты оны ерте анықтау асқынуларды уақтылы алдын алу үшін үлкен маңызға ие.

Зерттеудің мақсаты: негізгі өмір салты факторлары және әлеуметтік-демографиялық факторлары мен метаболикалық синдромның даму қаупі арасындағы байланысты зерттеу.

Зерттеудің материалдары мен әдістері. Бұл зерттеу Түркістан қаласындағы метаболикалық синдромды зерттеу жөніндегі «10 жылдық проспективті зерттеуде метаболикалық синдромы бар науқастардың тамақтану ерекшеліктерін, өмір салтын және комплаенттіліктің генетикалық аспектілерін зерттеу» (AP19676909) атты жобаның

бір бөлігі болып табылады. Іріктемеге Х.А. Ясауи атындағы Халықаралық қазақ-түрік университетінің емханасына тіркелген халықтан 552 адам қатысты. Зерттеуге шұғыл көмекті қажет ететін жедел жағдайлары, созылмалы ауыр декомпенсацияланған жағдайлары бар адамдар қатыспады.

Нәтижесі: Пациенттердің қорытынды іріктемесі 549 адамды құрады, себебі әлеуметтік-демографиялық факторлардың немесе липидтер мен көмірсулар алмасуының параметрлерінің бірі бойынша деректері жоқ қатысушылар статистикалық талдаудан алынып тасталды. Метаболикалық синдром және өмір салты көрсеткіштері арасындағы байланыс, соның ішінде жұмыс күні бойы отыру уақыты, физикалық белсенділік, көп өлшемді логистикалық регрессиялық талдау арқылы бағаланды. Метаболикалық синдром мен отыру уақыты арасындағы оң байланыс анықталды, ол жыныс, темекі шегу және алкогольді тұтыну сияқты потенциалды конфаундинг факторларға түзету жасағаннан кейін статистикалық маңыздылығын сақтап қалды. Керісінше, түзетілген логистикалық регрессиялық талдау нәтижесінде метаболикалық синдром және физикалық белсенділік деңгейі арасында статистикалық маңызды кері байланысы анықталды.

Қорытынды: Зерттеу нәтижесінде біздің іріктеме қатысушыларында метаболикалық синдром дамуында реттелетін қауіп факторлары (отырықшы өмір салты, физикалық белсенділік деңгейі) статистикалық маңызды рөлін көрсетті. Метаболикалық синдромы бар пациенттерде реттелетін өмір салты факторларын түзету арқылы жекелеңдірілген тактиканы ұстану асқынулардың алдын алуға көмектеседі.

Түйінді сөздер: метаболикалық синдром, физикалық белсенділік, өмір салты, отырықшы өмір салты.

For citation / Для цитирования / Дәйексөз үшін:

Turmanbayeva A.A., Sadykova K.Zh., Nuskabayeva G.O., Nurdinov N.S., Sarsenbayeva G.Zh., Kemelbekov K.S., Seidakhmetova A.A. The relationship between the main lifestyle factors and metabolic syndrome // *Nauka i Zdravookhranenie* [Science & Healthcare]. 2024. Vol.26 (5), pp. 47-54. doi 10.34689/SH.2024.26.5.006

Турманбаева А.А., Садькова К.Ж., Нускабаева Г.О., Нурдинов Н.С., Сарсенбаева Г.Ж., Кемельбеков К.С., Сейдахметова А.А. Взаимосвязь между основными факторами образа жизни и метаболическим синдромом // *Наука и Здравоохранение*. 2024. Т.26 (5). С. 47-54. doi 10.34689/SH.2024.26.5.006

Турманбаева А.А., Садькова К.Ж., Нускабаева Г.О., Нурдинов Н.С., Сарсенбаева Г.Ж., Кемельбеков К.С., Сейдахметова А.А. Өмір салтының негізгі факторлары мен зат алмасу синдромының өзара байланысы // *Ғылым және Денсаулық сақтау*. 2024. Т.26 (5). Б. 47-54. doi 10.34689/SH.2024.26.5.006

Introduction

Chronic non-communicable diseases are now a prevalent global health problem. According to the World Health Organization, non-communicable diseases, including cardiovascular disease (CVD), diabetes and obesity, are now responsible for about two thirds of deaths worldwide outcomes [9].

In recent years, metabolic syndrome (MetS) has become a major global health problem and has attracted much attention because it is strongly associated with various mortality[17].

Although different criteria for verification are used, the major components of MetS, including abdominal obesity, dyslipidemia, arterial hypertension, and insulin resistance, are to some extent related to sedentary lifestyle [6].

Along with the current trend of obesity worldwide, the rapid increase in the incidence and prevalence of MetS represents a major public health challenge that requires prompt and effective plans to address it [10].

Each component of the metabolic syndrome benefits to some extent from physical activity. Given that the prevalence of the metabolic syndrome and its individual components has increased significantly in recent decades, international guidelines call for greater efforts to reduce the incidence of this condition and its components. While physical activity that results in improved physical fitness cannot be expected to normalize insulin resistance, lipid abnormalities or obesity, the combined effect of a set of factors on these risk markers has been shown to have a

significant impact on health outcomes associated with the metabolic syndrome [4].

According to a study conducted between 2012 and 2014 in the Southern region of Kazakhstan, the prevalence of MetS was 23.9% (95% CI 18.6–29.2) among men and 21.8% (95% CI 18.5–25.2) among women. The authors of the study explained this finding by changes in diet and physical activity due to the fairly rapid economic development of Kazakhstan. They concluded that this high prevalence of MetS is alarming and calls for increased preventive measures. The authors also emphasized the need for further research into the risk factors that contribute to MetS in Kazakhstan, particularly lifestyle factors such as diet and physical activity. They suggested that this research could provide a basis for targeted interventions to reduce the prevalence of this condition. Thus, the problem of MetS does not lose its relevance, it requires a more detailed study of risk factors for the development of this pathology in each population. The study of the role of modifiable risk factors deserves special attention, as the main preventive measures should be directed at them. In view of the above, the aim of our study was to investigate the association between the main lifestyle and socio-demographic factors and MetS.

Material's and method: The present study is part of a large metabolic syndrome research project in Turkestan city "Investigation of genetic aspects of nutrition, lifestyle and compliance of patients with metabolic syndrome in a 10-year prospective study". (AP19676909). The sample

consisted of 552 patients from the attached population of the outpatient clinic of the H.A. Yasavi International Kazakh-Turkish University. Individuals with acute conditions requiring emergency care, chronic severe decompensated conditions were not included in the study. Patients lacking any of the anthropometric or clinical and metabolic parameters were excluded from statistical analysis. The final sample amounted to 549 patients.

The study included a questionnaire, anthropometric data and laboratory examination. For the purposes of this study, only the following information was used: sex, age, data on physical activity, smoking and alcohol consumption, as well as data on anthropometric and laboratory tests required for MetS determination. Anthropometric measurements were performed according to generally accepted requirements [2]. Waist circumference was (WC) measured with a soft centimeter tape, and the results were evaluated in centimeters (cm). WC was measured midway between the last ribs and the top of the pelvis at the upper anterior iliac crest (approximately at the level of the umbilicus). Blood pressure (BP) was measured using a non-automatic sphygmomanometer with the necessary requirements for correct BP recording. According to the results of two measurements with a 5-minute interval, the average BP was determined.

Blood sampling for laboratory analysis was performed in all patients from the ulnar vein after 12-hour fasting. All laboratory tests were performed in the laboratory of the clinic of the H.A. Yasavi International Kazakh-Turkish University with obligatory external and internal control.

Biochemical studies of blood lipid spectrum parameters included determination of total cholesterol (TC), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C) by standard enzymatic methods on a biochemical analyzer Cobas Integra-400 from Roche (Germany). Fasting blood glucose was determined after 12-hour fasting by taking capillary blood from the finger by express method using an Optium Xceed glucometer calibrated for venous blood plasma by Abbott Diabetes Care Ltd (UK). The results were evaluated in mmol/L.

MetS was diagnosed according to IDF (2005) criteria: abdominal obesity (waist circumference (WC) in men ≥ 94 cm, in women ≥ 80 cm) and any two of the following: 1) TG ≥ 1.7 mmol/L; 2) HDL-C in men < 1.03 mmol/L, in women < 1.29 mmol/L or hypolipidemic therapy; 3) CAD ≥ 130 or DA ≥ 85 mm Hg or antihypertensive therapy; 4) fasting glycemia ≥ 5.6 mmol/L or the fact of previously diagnosed type 2 DM [7].

Data were analyzed using the statistical software package SPSS 25 (SPSS Inc, Chicago, IL). Quantitative data were presented as mean values (M) and standard deviations (SD). Two independent samples were compared using Student's unpaired t-test. Categorical data were presented as percentages and analyzed using Pearson's chi-square test.

The association between the presence of MetS and lifestyle indicators, including sitting time during the working day and physical activity, was assessed using multivariate logistic regression analysis, in which MetS was used as a binary response variable. Independent variables were entered using the forced entry method. Gender, age,

smoking, and alcohol consumption were also entered into the model as potential confounding factors. Age was entered into the regression model as a categorical variable: <40 , 40-49, 50-59, and 60+ years. Smoking and alcohol use were entered as dichotomous variables. Unadjusted (uOR) and adjusted (aOR) odds ratios with 95% confidence intervals (CI) were calculated. For ranked independent variables, a test for linear trend was estimated by introducing the ranked variable categories into the model as a continuous variable. The critical level of significance (p) for statistical hypothesis testing was taken as 0.05.

The study was approved by the ethical committee of H. A. Yasavi International Kazakh-Turkish University (Protocol №30 dated 30.05.2024y.).

Results

The final patient sample was 549 patients, as study participants without data on any of the sociodemographic factors and on any of the lipid and carbohydrate metabolism parameters were excluded from the analysis.

Data from 126 men and 423 women were used for analysis. In the study population, smokers amounted to 10.0%. The proportion of alcohol drinkers amounted to 11.1%.

Socio-demographic and clinical characteristics of the study sample depending on gender are presented in Table 1. Statistically significant differences between men and women were revealed in age categories ($p=0.023$), smoking, and alcohol consumption. The proportion of MetS occurrence in the studied sample was statistically significantly ($p=0.003$) higher among men (52.4%) compared to women (37.1%). Both men and women showed comparable results in terms of time and level of physical activity (at least 150 min per week) (Table 1). The identified differences were taken into account in the subsequent regression analysis.

When comparing anthropometric and basic clinical and metabolic parameters depending on the sitting time during the working day, statistically significant differences were found in WC ($p=0.001$), thigh volume ($p<0.001$), body weight ($p<0.001$), triglyceride levels ($p=0.011$) and fasting glucose levels ($p<0.001$). The indicated parameters were higher in patients who spent 5 hours or more sitting during the day (Table 2).

To confirm the identified differences, a logistic regression analysis was conducted, taking into account potential confounding factors such as gender, age, smoking, and alcohol consumption. The results of the multivariate logistic regression analysis (Table 3) demonstrate a statistically significant, direct proportional relationship between sitting time and the presence of MetS, which was maintained after adjusting for confounding factors. (aOR=2.98, CI=2.01-4.42; p for trend <0.001). In contrast, a statistically significant inversely correlation was found with physical activity level after adjusted analysis (aOR =0.65, CI=0.43-0.88; p for trend=0.049). Statistically significant associations were found between the odds of having MetS according to gender (uOR=1.86, CI=1.25-2.78; p for trend=0.002), alcohol consumption (uOR =2.32, CI=1.35-3.98; p for trend=0.002), and smoking (uOR =1.87, CI=1.07-3.28; p for trend=0.028) in the unadjusted analysis.

Table 1.

Socio-demographic and clinical characteristics of the study sample.

		Total		Men (n=126)		Woman (n=423)		p
		n	%	n	%	N	%	
Age	<40	84	15,3	20	15,9	64	15,1	0,023
	40-49	134	24,4	26	20,6	108	25,5	
	50-59	166	30,2	29	23,0	137	32,4	
	> 60	165	30,1	51	40,5	114	27,0	
Alcohol Consumption	Yes	61	11,1	36	28,6	25	5,9	<0.001
	No	488	88,9	90	71,4	398	94,1	
Smoking	Yes	55	10,0	43	34,1	12	2,8	<0.001
	No	494	90,0	83	65,9	411	97,2	
Presence of MetS	Yes	223	40,6	66	52,4	157	37,1	0,003
	No	326	59,4	60	47,6	266	62,9	
Sitting time	Less than 5 hours during the working day	221	40,3	47	37,3	174	41,1	0,470
	5 hours or more during the working day	328	59,7	79	62,7	249	58,9	
Moderate-intensity physical activity (150 minutes per week or more)	No	407	74,1	87	69,0	320	75,7	0,164
	Yes	142	25,9	39	31,0	103	24,3	$\chi^2=2,207$ P=0,164

Table 2.

Anthropometric and basic clinical and metabolic parameters depending on sitting time.

Clinical and metabolic parameters	Sitting during the working day		p
	less than 5 hours a day	5 hours or more	
WC	93,31 (14,63)	97,39 (14,29)	0,001
Thigh volume	105,45 (11,30)	109,16 (12,15)	<0.001
Body weight	74,06 (14,67)	79,41 (16,24)	<0.001
SBP	122,84 (19,14)	124,42 (19,99)	0,352
DBP	79,60 (11,76)	80,06 (11,04)	0,642
TG	1,55 (0,72)	1,73 (0,97)	0,011
Fasting glucose	4,94 (0,55)	6,01 (2,23)	<0.001
HDL-C	1,36 (0,47)	1,28 (0,66)	0,130
LDL-C	2,74 (0,84)	3,02 (0,77)	0,272

After adjusting for the remaining potential confounding factors, the relationship between MetS and other variables such as gender, smoking, and alcohol consumption was not statistically significant. This suggests a significant role for

modifiable risk factors, such as a sedentary lifestyle and physical activity level, in the development of metabolic syndrome in our study population. Among non-modifiable risk factors, age remains statistically significant.

Table 3.

Binary logistic regression results: unadjusted (uOR) and adjusted (aOR) odds ratios with 95% confidence intervals (CI) in the study sample.

Indicator	MetS yes/no					
	uOR	95% CI	P	aOR *	95% CI	P
1	2	3	4	5	6	7
Sitting_time			<0,001			<0.001
Less than 5 hours during the working day	1,00	Reference		1,00	Reference	
5 hours or more per day during the working day	3,06	2,11; 4,44		2,98	2,01; 4,42	
Moderate-intensity physical activity (150 minutes per week or more)			0,128			0,049
No	1,00	Reference		1,00	Reference	
Yes	0,74	0,49; 1,09		0,65	0,43; 0,88	
Gender			0,002			0,156
Woman	1,00	Reference		1,00	Reference	
Men	1,86	1,25; 2,78		1,44	0,87; 2,38	
Age			<0,001			<0,001
<40	1,00	Reference		1,00	Reference	
40-49	2,24	1,12; 4,51		2,45	1,19; 5,05	
50-59	4,61	2,37; 8,97		4,74	2,38; 9,44	

Continuation of Table 3.

1	2	3	4	5	6	7
> 60	7,41	3,80; 14,44		7,15	3,57; 14,32	
Alcohol Consumption			0,002			
No	1,00	Reference		1,00	Reference	0,170
Yes	2,32	1,35; 3,98		1,56	0,84; 2,90	
Smoking			0,028			
No	1,00	Reference		1,00	Reference	0,177
Yes	1,87	1,07; 3,28		1,58	0,79; 3,19	
Note - * adjusted odds ratios were calculated with adjustment for age, gender, smoking and alcohol consumption						

Discussion

The results obtained from our study are similar to those of other similar studies. Lack of physical activity and poor cardiorespiratory fitness are consistently associated with an increased risk of chronic diseases, including type 2 diabetes and cardiovascular disease [13].

Exercise has well-described cardiovascular benefits and has been shown to improve biomarkers of MetS by reducing body weight, waist circumference and blood pressure, as well as improving circulating lipids and glucose profiles [8].

In combination with diet, physical activity is more effective in MetS compared to exercise alone or diet alone (67.4%, 23.5%, and 35.3%, respectively). The effect of a structured combined exercise program on weight loss is presumed to increase with prolonged exercise [11].

The Centers for Disease Control and Prevention (CDC) and American College of Sports Medicine (ACSM) guidelines recommend that all adults engage in at least 30 minutes of moderate-intensity physical activity per day to prevent chronic diseases, including type 2 diabetes and cardiovascular disease. These recommendations also emphasize the presence of physical activity, also its intensity and regularity. The findings suggest that men with sedentary lifestyles are more prone to metabolic syndrome than men who adhere to the CDC-ACSM recommendations, i.e., those who engage in at least 3 h·kg⁻¹ of at least moderate-intensity physical activity or physical activity. An active lifestyle controls low plasma high-density lipoprotein cholesterol (HDL-C) levels and reduces weight, hypertriglyceridemia, and arterial hypertension [19].

New evidence shows that reducing a sedentary lifestyle may also reduce metabolic risk. The American Diabetes Association and similar organizations in other countries have recently added recommendations for sedentary lifestyle to their Physical Activity Guidelines Activity Tracker-Based Metrics as Digital Markers of Cardiometabolic Health in Working Adults: Cross-Sectional Study [13].

A recent prospective cohort study involving 16,209 participants in Amsterdam, the Netherlands, showed that the prevalence of MetS in Dutch origin was 20.6% for men and 9% for women; whereas for participants of African Surinamese origin, the prevalence was 15.4% for men and 14.9% for women [15].

A cross-sectional study of 1,000 people aged 20 to 70 years living in an urban area in northern Iran found that high FA (compared with low FA) was inversely associated with MetS similar to a large study of 10,367 participants aged 37-66 years conducted in Poland. Also found that participants who reported low FA levels had a higher risk of

MetS than those who reported high FA levels. Moreover, a recent study involving 4865 adults in China found that higher levels of moderate and severe physical activity and total physical activity were associated with a lower risk of MetS [20].

A cross-sectional study of 750 patients from central rural India showed that physical activity was negatively correlated with TG and TC levels but did not correlate with HDL-CS levels [3].

Physical exercise significantly increased aerobic and muscular endurance, muscle mass and HDL. According to the results of a study conducted by Turkish scientists prolonged sitting time is associated with impaired lipid metabolism, while sufficient physical activity levels are favorably associated with fasting fat oxidation metabolic flexibility in sedentary and physically inactive adults with MetS. Although a causal relationship could not be determined due to cross-sectional study conditions, the results suggest that reducing sedentary lifestyle time and increasing physical activity of even mild intensity may help in the prevention of metabolic diseases in at-risk groups due to their favorable effects on lipid and carbohydrate metabolism [14].

The large geographic variation in the prevalence of MetS emphasizes the importance of environmental and lifestyle factors, such as excess calories in the diet and physical inactivity, as major contributors to the disease [1]. Low physical fitness is a major risk factor for MetS and overall mortality. Convincing evidence has shown that a high amount of sedentary behavior is associated with an increased risk of developing a range of chronic diseases and mortality [12].

The WHO recommendation is to limit time in a sedentary position in favor of even low-intensity activity. More health benefits can be gained by being active for at least 300 minutes per week [19]. Each clinical component of MetS can be modified by physical activity, which is an inexpensive and therefore readily available way to prevent and treat metabolic syndrome [18].

Conclusion.

The results of the conducted research suggest a statistically significant role for modifiable lifestyle factors, such as a sedentary lifestyle and lack of physical activity, in the development of metabolic syndrome. Personalized management tactics for patients with metabolic syndrome, by correcting modifiable lifestyle factors, can help prevent complications. To confirm these causal relationships, it is necessary to conduct further research. Specifically, it would be useful to analyze the influence of these lifestyle factors

on changes in clinical and metabolic parameters within a cohort over time, as part of a prospective study.

Competing interests: None declared.

Acknowledgements: None.

Funding: This research is funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant no. AP19676909)

Authors' contributions:

Nuskabayeva G.O. and Sadykova K.Zh. conceived the study; Turmanbayeva A.A. and Nurdinov N.S. acquired the data; Sadykova K.Zh., Kemelbekov K.S. and Nuskabayeva G.O. contributed to data analysis;

Turmanbayeva A.A., Sarsenbayeva G.Zh. and Sadykova K.Zh. drafted the manuscript;

Nuskabayeva G.O., Seidakhmetova A.A. and Nurdinov N.S. contributed to critical evaluation and revisions of the manuscript.

All authors read and approved the final version of the paper.

References:

1. Alsten S.K., Rabkin K.S., Sawada N., Shimazu T., Charvat H., Yamaji T., et al. Metabolic syndrome, physical activity, and inflammation: a cross-sectional analysis of 110 circulating biomarkers in Japanese adults. *Cancer Epidemiol. Biomarkers Prev.* 2020. (8), 1639-1646. www.pub.med.com.29 http://doi.org.10.1158/1055-9965.Epi-19-1513.

2. Assel Sadykova, Zhanar Shalkharova, Zhanat Shalkharova, Karlygash Sadykova, Kamshat Madenbay, Mira Zhunissova, Gulnaz Nuskabayeva, Saltanat Askarova, Andrej Grijbovski Metabolic syndrome and its components in southern Kazakhstan: a cross-sectional study», *International Health*, Volume 10. Lancet, Issue 4, July 2018, Pages 268–276, https://doi.org/10.1093/inthealth/ihy 027.

3. Babita Bondge, Jyoti Jain, Mangesh Warkad, Madhura Joshi, Subhash More, Saiprasath Janaarthan. Association of Physical Activity with Lipid Profile in Healthy Subjects: A Cross Sectional Study in Tertiary Care Hospital from Central Rural India. www.pub.med.com. 2021 25(6): 520-526. Nov-Dec. http://doi.org.10.4103/ijem.ijem_327_21.

4. Cuchel M., McGowan M.P. Familial hypercholesterolaemia: too many lost opportunities. 2021 Nov 6. 398(10312): 1667-1668. www.pub.med.com. http://doi.org/10.1016/S0140-6736(21)01372-6.

5. Catalina Mascaró, Cristina Bouzas, Josep Tur. Association between Non-Alcoholic Fatty Liver Disease and Mediterranean Lifestyle: A Systematic Review. 2021 23. 14(1):49. Dec. www.pub.med.com http://doi.org/10.3390/nu14010049.

6. Esther M.F. van Sluijs, Ulf Ekelund, Inacio Crochemore-Silva et al. Physical activity behaviours in adolescence: current evidence and opportunities for intervention. 398(10298): 429-442. Affiliations expand, Jul 31. http://doi.org/10.1016/S0140-6736(21)01259-92021.

7. International Diabetes Federation. The IDF consensus worldwide definition of the metabolic syndrome. Available at: http://www.idf.org/webdata/docs/IDF_Metasyndrome_definition.pdf (accessed 24.08.2005).

8. Irene Sequi-Dominguez, Celia Alvarez-Bueno, Vicente Martinez-Vizcaino, Rubén Fernandez-Rodriguez,

Alicia Del Saz Lara, Iván Caverro-Redondo. Effectiveness of Mobile Health Interventions Promoting Physical Activity and Lifestyle Interventions to Reduce Cardiovascular Risk Among Individuals With Metabolic Syndrome: Systematic Review and Meta-Analysis. 31. 22(8):e17790. www.pub.med.com. 2020 Aug http://doi.org/10.2196/17790.

9. Jonathan Myers, Peter Kokkinos, Eric Nyelin. Physical Activity, Cardiorespiratory Fitness, and the Metabolic Syndrome. 2019 Jul; 11(7): 1652. www.pub.med.com. http://doi.org/10.3390/nu11071652.

10. Jonathan Myers, Peter Kokkinos, Eric Nyelin. Physical Activity, Cardiorespiratory Fitness, and the Metabolic Syndrome». 2019 Jul 19. www.pub.med.com. PMID:31331009. http://doi.org/10.3390/nu11071652PMCID: PMC6683051.

11. Joseph M.S., Tincopa M.A., Walden P., Jackson E., Conte M.L., Rubenfire M. The Impact of Structured Exercise Programs on Metabolic Syndrome and Its Components: A Systematic Review». *Diabetes Metab. Syndr. Obes.* 2019. 12: 2395–2404. http://doi.org/10.2147/DMSO.S211776.

12. Koenigstein K., Dipla K., Zafeiridis A. Vascular exercise: molecular and clinical effects of exercise on vascular health. A narrative review. *Cells* 2023. 12 (21), 2544. http://doi.org.10.3390/cells12212544.

13. Lakka T.A., Laaksonen D.E., Lakka H.-M., Mannikko N., Niskanen L.K., Rauramaa R. Sedentary Lifestyle, Poor Cardiorespiratory Fitness, and the Metabolic Syndrome. *Med. Sci. Sports Exerc.*, Vol. 35, No. 8, pp. 1279–1286, 2019. http://doi.org/10.2196/17798.

14. Long Y., Tang L., Zhou Y., Zhao S., Zhu H. Causal relationship between gut microbiota and cancers: a two-sample Mendelian randomisation study. *BMC Med.* 2023 Feb 21. 21(1):66. http://doi.org.10.1186/s12916-023-02761-6.

15. Manon Balvers, Marcus de Goffau, Natal van Riel, Bert-Jan van den Born et al. Ethnic variations in metabolic syndrome components and their associations with the gut microbiota: the HELIUS study. 2024. 20. 16(1):41. Mar. www.pub.med.com. http://doi.org/10.1186/s13073-024-01295-7.

16. Oppert D.M., Bellich A., Chiangur K. Physical activity in the management of obese individuals. *Eur. J. Intern Med.* 2021. 93, 8-12. http://doi.org.10.1016/j.ejim.2021.04.028

17. Saklayen M.G. The global epidemic of the metabolic syndrome. *Curr Hypertens Rep.* 2018. 20:12. http://doi.org/10.1007/s11906-018-0812-z.

18. Tomasz Chomiuk, Natalia Niezgodna, Natalia Niezgodna, Artur Mamcarz, Artur Mamcarz, Daniel li, Daniel Śliż. *Physiol. Exercise Physiology* Volume 19 February 2024 Sec. http://doi.org.10.3389/fphys.2024.1365761.

19. Yuri Rykov, Thuan-Quoc Thach, Gerard Dunleavy, et al. Activity Tracker-Based Metrics as Digital Markers of Cardiometabolic Health in Working Adults: Cross-Sectional Study. 2020 Jan 31;8(1):e16409. www.pub.med.com. http://doi.org/10.2196/16409.

20. Yuxuan Tan, Yingyin Fu, Huojie Yao, Xiaomei Wu, Zhiyu Yang et al. Relationship between phthalates exposures and hyperuricemia in U.S. general population, a multi-cycle study of CHINES 2007-2016. 2023 Feb 10. 859(Pt.1): 160208. www.pub.med.com. http://doi.org/10.1016.

Information about authors:

Ainur A. Turmanbayeva - Master of medical sciences, 2nd year doctoral student on the educational program D141 Medicine (8D10110-«Medicine»), ORCID 0009-0004-4263-8157, Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkistan city, Republic of Kazakhstan. Postal address: 161200, Republic of Kazakhstan, Turkestan, Bekzat Sattarkhanov Avenue, 29/3, E-mail: ainur.turmanbaeva@ayu.edu.kz; Phone: +7 747 816 93 37.

Karlygash Zh. Sadykova - PhD, associate professor, head of «Special clinic subjects» department, ORCID 0000-0002-9120-8565; Khoja Akhmet Yassawi International Kazakh-Turkish University; Turkistan city, Republic of Kazakhstan; e-mail: karlygash.sadykova@ayu.edu.kz, Phone: +7 707 731 64 76

Gulnaz O. Nuskabaeva - Candidate of medical sciences, associate professor, head of «Special clinic subjects» department, ORCID 0000-0003-2139-3221; Khoja Akhmet Yassawi International Kazakh-Turkish University; Turkistan city, Republic of Kazakhstan; e-mail: nuskabaeva.gulnaz@ayu.edu.kz, Phone: +7 705 285 31 31

Nursultan S. Nurdinov - PhD, senior lecturer, department of "Special clinic subjects", ORCID 0000-0001-5341-7211; International Kazakh-Turkish University named after H.A. Yassavi; Turkistan city, Republic of Kazakhstan; e-mail: nursultan.nurdinov@ayu.edu.kz; Phone: +7 705 378 78 91

Gulzat Zh. Sarsenbayeva – Candidate of Medical Sciences, Associate Professor, Head of the Department "Social health insurance and public health", num: 87017496288, e-mail: gulzat.zh@mail.ru; <https://orcid.org/0000-0002-1518-6528>, JSC South Kazakhstan Medical Academy, Shymkent, Republic of Kazakhstan;

Kanatghan S. Kemelbekov – PhD., Head of the Department "Pediatrics-1, num: 87085334300, e-mail: kanat-270184@mail.ru; <https://orcid.org/0009-0008-1619-6278>, JSC South Kazakhstan Medical Academy, Shymkent, Republic of Kazakhstan;

Aizat A. Seidakhmetova – Candidate of Medical Sciences, Head of the Department of "Emergency Medicine and Nursing", num: 87017671448, e-mail: aizat67@mail.ru; <https://orcid.org/0000-0002-5794-3814>, JSC South Kazakhstan Medical Academy, Shymkent, Republic of Kazakhstan;

Corresponding author:

Turmanbayeva Ainur Azimkhanovna - Master of medical sciences, 2nd year doctoral student on the educational program D141 Medicine (8D10110-«Medicine»), ORCID 0009-0004-4263-8157, Khoja Akhmet Yassawi International Kazakh-Turkish University, Turkistan city, Republic of Kazakhstan.

Postal address: 161200, Republic of Kazakhstan, Turkestan, Bekzat Sattarkhanov Avenue, 29/3

E-mail: ainur.turmanbaeva@ayu.edu.kz.

Phone: +7 747 816 93 37